# Traditional Knowledge and Useful Plant Richness in the Tehuacán–Cuicatlán Valley, Mexico[1]

Rafael Lira*[,2], Alejandro Casas[3], Rocío Rosas-López[2], Martín Paredes-Flores[2], Edgar Pérez-Negrón[3], Selene Rangel-Landa[3], Leonor Solís[3], Ignacio Torres[3], and Patricia Dávila[2]

[2]Laboratorio de Recursos Naturales UBIPRO, Facultad de Estudios Superiores Iztacala, Universidad Nacional Autónoma de México, Mexico City, Mexico
[3]Centro de Investigaciones en Ecosistemas, Universidad Nacional Autónoma de México, Mexico City, Mexico
*Corresponding author; e-mail: rlira@servidor.unam.mx

**Traditional Knowledge and Useful Plant Richness in the Tehuacán–Cuicatlán Valley, Mexico.** This study systematizes ethnobotanical information about the interactions between people and plants, ethnofloristic richness, the relative importance of useful species richness in relation to general species richness, and plant management in the Tehuacán–Cuicatlán Valley of central Mexico. The study recorded a total of 1,605 useful vascular plant species (61.2% of the total species richness of the regional vascular flora), this being the region with the highest absolute richness of useful plant species in Mexico. The null hypothesis that plant families with a higher number of useful species would be those having a higher general species richness was analyzed through residuals method. The plant families richest in useful species were Poaceae, Asteraceae, Cactaceae, Cyperaceae, Mimosaceae, and Solanaceae, most of which also have the highest general floristic richness. However, analyses of use categories did not generally corroborate our hypothesis. About 1,335 of the useful species are wild, more than 500 species are submitted to some type of management (62 species are tolerated, 34 protected, 50 enhanced, and 358 cultivated), but only a few have been studied to document their process of domestication. This information can be useful for developing regional strategies of sustainable management of plant resources.

**Conocimiento tradicional y riqueza de plantas útiles en el Valle de Tehuacán-Cuicatlán, México.** Este trabajo sistematiza información etnobotánica sobre las interacciones entre la gente y las plantas, la riqueza etnoflorística, la importancia relativa de las especies útiles con respecto a la riqueza florística general, y sobre el manejo de plantas en el Valle de Tehuacán–Cuicatlán, en el centro de México. Se registró un total de 1,605 especies de plantas vasculares útiles (61.2% de la riqueza total de la flora vascular de la región), lo que identifica a la región como la de mayor riqueza de plantas útiles de México en términos absolutos. Mediante el método de residuales se analizó la hipótesis nula de que las familias con mayor número de especies útiles serían las de mayor riqueza florística. Se encontró que las familias con mayor número de especies útiles fueron Poaceae, Asteraceae, Cactaceae, Cyperaceae, Mimosaceae, y Solanaceae, las cuales en su mayoría son las de mayor riqueza florística en la región. No obstante, al efectuar el análisis por categorías de uso no siempre se confirmó nuestra hipótesis. Casi 1,335 de las plantas útiles son silvestres, pero alrededor de 500 están sometidas a algún tipo de manejo (62 especies son toleradas, 34 protegidas, 50 fomentadas y 358 cultivadas), pero sólo para algunas de ellas se ha estudiado cómo operan los proceso de domesticación. La información generada en el trabajo puede ser útil para el desarrollo de estrategias regionales de manejo sustentable de los recursos vegetales.

**Key Words:** Ethnobotany, Tehuacán–Cuicatlán Valley, traditional plant management, residual analysis, México.

## Introduction

The cultural area known as Mesoamerica, between central Mexico and northern Costa Rica, is currently recognized as one of the most important centers of cultural diversity and biological richness of the world (Hernández-Xolocotzi 1993; Rzedowski 1993; Toledo et al. 1997). Humans and plants have interacted here for nearly 12,000 years (MacNeish 1992). Local Mesoamerican peoples have developed

[1] Received 2 July 2008; accepted 4 February 2009; published online _____.

*Economic Botany*, XX(X), 2009, pp. 1–17.
© 2009, by The New York Botanical Garden Press, Bronx, NY 10458-5126 U.S.A.

Published online: 05 March 2009

an extraordinary complex of interaction with plants, including the gathering of wild plants, several forms of *in situ* or silvicultural management of wild plant populations and communities, and the cultivation and selection of particular varieties to be used in specific environmental and cultural conditions (Caballero 1984; Bye 1993; Casas et al. 1997a, 2007).

Such a long and systematic interaction has led to the construction of rich knowledge profiles regarding the use and management of a wide range of plant species. It has been estimated that nearly 5,000 to 7,000 plant species are used in Mexico, including their immense infra–specific variation (Caballero 1984; Casas et al. 1994). Nearly 90% of the useful plant species are native wild species obtained through gathering (Caballero et al. 1998), whereas nearly 20 plant species (such as corn, beans, cocoa, chili peppers, cotton, and squashes) are important economic crops on a global scale and whose extraordinary diversity of land races and populations of wild relatives constitute highly relevant genetic resources. Some other native Mexican species (nearly 200 species, according to Casas and Parra 2007) are in the early and intermediate stages of domestication. All of them and their wild relatives are not only important on a local or regional scale, but also represent a bulk of potential plant resources that have not been well studied, although they have already been documented by various ethnobotanical studies, such as those reported by Alcorn (1984), Bye (1993), Caballero and Mapes (1985), Caballero et al. (2000), Casas and Caballero (1996), Casas et al. (1996, 1997a, 1999a, 2001, 2007), Lira (2004), Lira and Caballero (2002), Lira and Casas (1998), Martínez-Alfaro et al. (1995), and Zizumbo-Villarreal and Colunga-García-Marín (1982).

The Tehuacán–Cuicatlán Valley is one of the Mexican areas where the relationships between humans and plants have been relevant in the past and still are today. In this region, the interactions between local people and plants from prehistory to the present day have been documented (MacNeish 1967; Smith 1967; Casas et al. 2001). However, until the end of the last century, ethnobotanical studies were relatively scarce in this area. Miranda (1948), Smith (1965, 1967), Casas and Valiente-Banuet (1995), and Casas et al. (1997a) conducted some of the most relevant work.

Casas et al. (2001) synthesized the ethnobotanical information generated in the region until the beginning of the present century. It included an inventory of common names, uses, and management levels for 808 species of vascular plants from seven municipalities of the region, as well as an analysis of the ethnofloristic importance of the Tehuacán Valley compared with other regions of Mexico. The result of this analysis highlighted the valley as one of the most important regions of Mexico in terms of richness of traditional knowledge of plants. In the last few years, however, our group has conducted a series of new studies in the area, which have contributed to the depth of the ethnobotanical framework and to the expansion of the spatial, ecological, and cultural limits covered in the previous study (Casas et al. 2001). The new data practically double the number of useful flora reported by those authors. Therefore, new analyses of the information are necessary considering the outstanding richness of ethnobotanical knowledge documented and considering the importance of the region in the study of the history of interactions between humans and plants that led to the origins of agriculture.

Accordingly, this study aimed to report and systematize the ethnobotanical investigations carried out to date. Particular attention was given to information on the regional ethnofloristic richness, since the Tehuacán–Cuicatlán Valley is the area of Mexico with the richest useful flora reported hitherto. Also, an analysis was undertaken to evaluate the relative importance of plant families in relation to their richness of useful plant species and their general floristic richness in the region. And finally, information on various forms of plant management occurring in the region were also analyzed in the context of one of the most important scenarios of domestication of plants recognized in Mexico by archaeological and ethnobotanical studies.

## Study Area

The Tehuacán–Cuicatlán Valley constitutes a floristic province belonging to the Mexican Xerophytic Region (Rzedowski 1978). It is located in the southern part of the state of Puebla and the northern portion of the state of Oaxaca (Fig. 1). The surface of the valley is about 10,000 square kilometers ($km^2$), the region forming a complex physiographic mosaic with several internal valleys separated by mountain chains. The climate is predominantly semiarid, with an annual mean temperature of 21°C and an annual mean




**Fig. 1.** Geographic position of the Tehuacán–Cuicatlán Valley and the localities where the ethnobotanical information was gathered.

precipitation of 400 millimeters (mm) (Villaseñor et al. 1990). The dryness of the region is the result of the orographic rain shadow effect produced by the Oriental Sierra Madre (Villaseñor et al. 1990; Valiente-Banuet 1991; Dávila et al. 2002). The altitude of the floor of the valley ranges from 2,200 meters (m) in the northwest to about 600 m in the southeast; the dryland area is below 1,800 m. This region has been recognized as a center of megadiversity and endemism by the World Conservation Union (IUCN) and, in 1998, most of its territory was decreed as part of the Tehuacán–Cuicatlán Biosphere Reserve (Dávila et al. 1993, 2002; Mendez-Larios et al. 2005). Its vascular flora comprises 2,621 species and 48 infra–specific taxa of 180 families and 890 genera, with between 9.1% and 13.9% of the plant species being endemic to the region (Dávila et al. 1993, 2002; Mendez-Larios et al. 2004, 2005). At the plant community level, Valiente-Banuet et al. (2000) have identified 30 different plant association types.

The biological richness recorded at this point in time includes 57 species of mosses (Delgadillo and Zander 1984), 34 species of bats (Rojas-Martínez and Valiente-Banuet 1996), 141 species of birds (Arizmendi and Espinoza de los Monteros 1996), 24 species of Coreidae (Hemiptera–Heteroptera) (Brailovsky et al. 1994, 1995), and 24 species of ants (Ríos-Casanova et al. 2004), among other groups. Compared with other areas, these figures reveal that the Tehuacán–Cuicatlán Valley is probably the most diverse arid region of North America (Dávila et al. 2002) and it is now an important natural protected area of Mexico.

The Tehuacán–Cuicatlán Valley also has a high cultural diversity. Of the 57 indigenous ethnic groups that currently inhabit the Mexican territory (Toledo et al. 2001), seven (Nahua, Popoloca, Mazatec, Chinantec, Ixcatec, Cuicatec,

and Mixtec) live in the valley. The region has been important in the reconstruction of the prehistory of Mesoamerica since archaeologists found in caves some the oldest remains of domesticated plants and agriculture in the New World (MacNeish 1967, 1992; Smith 1967). At present, the Tehuacán–Cuicatlán Valley is densely populated and its inhabitants undertake numerous economic activities, especially agriculture and livestock farming. However, there are areas with various levels of degradation and others that are well preserved, such as where people still practice the gathering of animal and plant products to complement their diet and daily subsistence (Casas et al. 2001; Pérez-Negrón and Casas 2007).

## Methods

### Ethnobotanical Surveys

Recent ethnobotanical studies were focused on undertaking inventories of the plant species used by local peoples, the indigenous nomenclature and forms of management, collecting and georeferencing herbarium specimens, and evaluating the use rates and spatial availability of plant resources. The studies comprised the villages of (1) Santa María Ixcatlán, in the municipality of Ixcatlán (Oaxaca), which is inhabited by Ixcatec people (the only village of this indigenous ethnic group in the World) (Rangel-Landa and Lemus 2002), (2) the village of San Rafael, in the municipality of Coxcatlán (Puebla) where mestizo people with Náhuatl cultural influence live (Rosas-López 2003; Blanckaert et al. 2004), (3) the villages of San Pedro Nodón and San Pedro Jocotipac, in the municipalities of Cuicatlán and Jocotipac (Oaxaca), respectively, which are inhabited by Mixtec people (Echeverría-Ayala 2003); (4) the village of San Luis Atolotitlán, in the municipality of Caltepec (Puebla), which is inhabited by mestizo people with Náhuatl cultural influence (Torres 2004); (5) the village of San Lorenzo Pápalo, in the municipality of Cuicatlán (Oaxaca) with the Cuicatec people, one of the endemic ethnic groups of the Tehuacán Valley (Solís 2006); (6) the village of Zapotitlán de las Salinas, in the municipality of Zapotitlán (Puebla), which is inhabited by mestizo people with Popoloca origins (Paredes-Flores et al. 2007); and (7) the village of Quiotepec, in the municipality of Cuicatlán (Oaxaca), with mestizo people (Pérez-Negrón and Casas 2007). Table 1 summarizes the information related to these study areas, and Fig. 1 shows their geographic localization in the Tehuacán–Cuicatlán Valley.

We determined the ethnofloristic richness of the Tehuacán–Cuicatlán Valley by systematizing in a database the information of all these case studies, including the data previously synthesized by Casas et al. (2001), and ethnobotanical information documented in studies on the domestication of columnar cacti (Casas et al. 1997b, 1999a, 1999b, 2006; Cruz and Casas 2002; Arellano and Casas 2003; Carmona and Casas 2005; Rodríguez-Arévalo et al. 2006; Blancas 2007; Parra et al. 2008), as well as other species such as *Sideroxylon palmeri* (González-Soberanis and Casas 2004), *Proboscidea louisianica* ssp. *fragans* (Paredes-Flores 2006), and *Ceiba aesculifolia* ssp. *parvifolia* (Avendaño et al. 2006).

### General Floristic Richness and Richness of Resources of Plant Families

Our group analyzed the relative importance of plant families, in terms of general species richness and richness of useful plant species, through a residual analysis using the method developed by Moerman et al. (1999) and recently used by Caballero et al. (2004). Through this method, a regression of the total number of useful species and the number of species per use category per plant family, reported in the ethnobotanical studies referred above, was performed with respect to the total number of species per plant family reported for the region by Dávila et al. (1993, 2002) and Mendez-Larios et al. (2005). Databases constructed by those authors were compared with ours. After the regression was performed, the families were ordered hierarchically according to their residual values, which represent the difference between the predicted number of useful species obtained by the regression and the real number of species documented by the ethnobotanical studies. Accordingly, the families with higher positive residual values are those that are more favored as useful plants and/or for any particular use. In contrast, the families with negative residual values are those less frequently used or even ignored by people. Thus, this method was used to test the null hypothesis that if no use selectivity is practiced on particular groups of plant resources, the best–represented families in the flora will be those having higher number of useful species.

**Table 1.** General characteristics of the localities where the ethnobotanical information was gathered in the Tehuacán–Cuicatlán Valley.

| Sites or Regions | Altitude (in meters) | Vegetation | Ethnofloristic Richness | Human Group | References |
|---|---|---|---|---|---|
| Municipalities of Zapotitlán Salinas (San Juan Raya, Los Reyes Metzontla), Tehuacán (San Lorenzo, Coapan, Tehuacán), Coxcatlán and Ajalpan (Ajalpan, San Rafael, Rancho El Aguaje, Coxcatlán), Cuicatlán (Cuicatlán, Tomellin, Valerio Trujano), Chazumba (Chazumba), and San Pedro and San Pablo Tequixtepec (Tequixtepec, Santa Catalina Chinango) | 1,000–2,300 | Thorn–scrub forest, tropical deciduous forest, and Mexical or Chaparral | 808 | Chocho–Popoloca, Nahua, Cuicatec, Mixtec, and Mestizo | Casas et al. (2001) |
| Zapotitlán Salinas | 1,400–1,600 | Thorn–scrub forest | 288 | Chocho–Popoloca and Mestizo | Paredes-Flores (2006), Paredes-Flores et al. (2007) |
| Santa María Ixcatlán | 1,840 | *Quercus* forest, *Juniperus flaccida* forest, riparian forest with *Taxodium mucronatum,* rosette scrub forest, or Izotal with *Beaucarnea gracilis*, Mexical or Chaparral, Palmar with *Brahea dulcis*. | 376 | Ixcateco | Rangel-Landa and Lemus (2002) |
| San Rafael | 1,217 | Thorn–scrub forest | 368 | Nahua and Mestizo | Blanckaert et al. (2004), Rosas-López (2003) |
| San Pedro Nodón and San Pedro Jocotipac | 1,680 | Cardonal with *Mitrocereus fulviceps,* or *Cephalocereus columna–trajani, Quercus* forest, Palmar with *Brahea dulcis*, riparian forest with *Taxodium mucronatum*. | 264 | Mixteco | Echeverría-Ayala (2003) |
| Santiago Quiotepec | 545 | Jiotillal with *Escontria chiotilla*, Tetechera with *Neobuxbaumia tetetzo*, Cardonal with *Pachycereus weberi*, riparian forest with *Salix chilensis* | 252 | Mestizo | Pérez-Negrón and Casas (2007) |

**Table 1.** (Continued).

| Sites or Regions | Altitude (in meters) | Vegetation | Ethnofloristic Richness | Human Group | References |
|---|---|---|---|---|---|
| San Lorenzo Pápalo | 1,800 | Deciduous tropical forest, *Quercus* and *Quercus–Pinus* forest, *Alnus* forest, and riparian forest. | 367 | Cuicateco | Solís (2006) |
| San Luis Atolotitlán | 1,900 | Chichipera, Cardonal dominated by *Mitrocereus fulviceps*, Izotal with *Beaucarnea purpusii*, rossette–scrub forest with *Dasylirion serratifolium*, *Quercus* forest, Palmar with *Brahea dulcis*. | 280 | Mestizo | Torres (2004) |

# Results

## The Useful Flora of the Tehuacán–Cuicatlán Valley

The ethnofloristic inventory reported a total of 1,605 useful vascular plant species belonging to 671 genera and 147 families. The entire inventory is available as an electronic Appendix of this publication and can be consulted by contacting the authors. The comparison of these ciphers with those reported for other areas of Mexico highlights the importance of the Tehuacán Valley as the region with the highest absolute richness of useful plant species and one of the highest relative richness in relation to other areas, second only to the Sierra de Manantlán, Jalisco (Table 2). The inventory identified a total of 33 use categories of plant species, which contribute to satisfy a broad spectrum of human needs (Table 3). Based on the number of species belonging to each use category, the most outstanding uses are fodder, medicinal, food, firewood, ornamental, as well as wood and construction material.

## Relative Importance of the Plant Groups

The most important plant families according to their general species richness in the region are Poaceae, Asteraceae, Fabaceae, Cactaceae, and Mimosaceae (Table 4), which are also important physiognomic elements of the plant communities of the area. The residual values revealed that the richest plant families in useful species are Poaceae, Asteraceae, Cactaceae, Cyperaceae, Mimosaceae, and Solanaceae; as mentioned, most of them have a high general floristic richness. However, in the valley there are some other families that also have high floristic richness, such as Fabaceae and Euphorbiaceae; they are not as proportionally important as sources of useful plants (Table 5).

Analyzing the data according to the use categories, the results indicate that certain families are sources of specific resources (see Table 6). For instance, among the plants that are used as fodder, the Poaceae and Cyperaceae are the most demanded groups, and practically all of their species are utilized for this purpose. In contrast, other families with high floristic richness, such as Asteraceae, Fabaceae, Euphorbiaceae, and Solanaceae, are the least used in this category. Among the ornamental plants, three of the most diverse plant groups (Asteraceae, Cactaceae, and Solanaceae) and three of the least rich (Crassulaceae,

**Table 2.** Relative richness of the useful species of the Tehuacán–Cuicatlán Valley, compared to other regions of Mexico.

| Region | Useful Species | Area (km$^2$) | Relative Richness[1] | Reference |
|---|---|---|---|---|
| Valle de Tehuacán, Oaxaca and Puebla | 1,605 | 10,000 | 0.16 | This study |
| Sierra Norte de Puebla, Puebla | 720 | 13,000 | 0.06 | Martínez-Alfaro et al. (1995) |
| Uxpanapa, Veracruz | 325 | 5,000 | 0.07 | Toledo et al. (1995) |
| Selva Lacandona, Chiapas | 415 | 13,000 | 0.03 | Toledo et al. (1995) |
| Sian Ka'an, Quintana Roo | 316 | 5,280 | 0.06 | Toledo et al. (1995) |
| Huasteca, San Luis Potosí | 445 | – | – | Alcorn (1984) |
| Tepehuan region, Chihuahua | 380 | 10,500 | 0.04 | Pennington (1969) |
| Sierra de Manantlán, Jalisco | 650 | 1,400 | 0.46 | Benz et al. (1994) |
| Tzeltal region of Tenejapa, Chiapas | 645 | – | – | Berlin et al. (1974) |
| Península de Yucatán | 1,000 | 140,056 | 0.01 | Flores (1999) |
| Region de La Montaña, Guerrero | 430 | 11,000 | 0.04 | Casas et al. (1994) |

[1] Estimated as a ratio among useful species and area of the region.

Convolvulaceae, and Malvaceae) showed high residual values. We found contrasting results in the case of species used as wood and construction material, among which Fagaceae and Mimosaceae had higher residual values than Asteraceae and Poaceae, which are two of the richest plant families in the valley.

The analysis allows appreciation of the relevant sources of human food plant families that are relatively well represented in the region, such as Cactaceae, Mimosaceae, Fabaceae, and Solanaceae. This is also the case of Cucurbitaceae, which, although is not one of the richest families of the region, it is recognized as an important source of edible plants. The highest residual values of species used as firewood were recorded for Mimosaceae and Fagaceae, whereas other floristically richer families had lower (Asteraceae and Cactaceae) or negative residual values (Poaceae and Euphorbiaceae). Finally, among the medicinal plants, Asteraceae is the family with the highest residual value, followed by other 10 families with markedly lower residual values. It is important to note that these families constitute a mixture of plant groups including well–represented families in the Tehuacán–Cuicatlán flora, such as Euphorbiaceae and Lamiaceae, as well as other less diverse families, such as Burseraceae, Fagaceae, and Cucurbitaceae. Other rich plant families of the region, such as Poaceae, Fabaceae, and Cactaceae, had low residual values, indicating that they are not primary sources of medicinal plants.

### Use and Management of Plant Resources

A total of 1,011 species had a unique category of use, whereas the remaining 594 (37%) had two or, in the case of 147 species, more than four use categories (Table 7). Ethnobotanical studies reveal that the local people in the region actively manage their plant resources, using practices directed to maintain their availability or even increasing their amounts and/or improving their quality. Nearly 1,335 of the 1,605 useful species are gathered in the wild, but many of them are also submitted to some of the following forms of management: (1) Selective tolerance (62 spp.), through which people deliberately leave individual plants of particular species and sometimes particular phenotypes of these species in sites where the ecosystem is being perturbed; (2) protection (34 spp.), through which people undertake periodic prunings, remove plants that compete with them, protect them from herbivores, frosting, and other factors; (3) enhancement (40 spp. by transplanting vegetative parts and 10 through their sexual propagules), through which people promote the dispersion of sexual and vegetative propagules to expand their availability and/or phenotype transplantation (people remove some complete individuals from their natural site to other locations), or even the cultivation of the latter ones. Finally, 358 species are cultivated in different agricultural systems, with the highest diversity found in home gardens. Examples of plant species submitted to one or more management forms are presented in Table 8.

## Discussion

To date, the richness of useful plant species recorded in the Tehuacán Valley represents

**Table 3.** Categories of use of the useful flora of the Tehuacán–Cuicatlán Valley and the species number represented in each category.

| Uses | Number of Species |
|---|---|
| Fodder | 874 |
| Medicinal | 396 |
| Food | 339 |
| Ornamental | 313 |
| Firewood | 209 |
| Wood and Construction Materials | 129 |
| Living Fences | 53 |
| Soils Control | 51 |
| Ceremonial | 50 |
| Handicraft | 47 |
| Shade | 41 |
| Toxic / Poisons | 26 |
| Resins / Latex | 22 |
| Domestic Tools | 17 |
| Fibers | 17 |
| Sources of Saponins | 15 |
| Alcoholic Beverages | 14 |
| Glues | 12 |
| Dyes | 9 |
| Commercial | 9 |
| Spices | 9 |
| Stimulants | 9 |
| Toys | 8 |
| Insecticide | 7 |
| Aromatizing | 4 |
| Edible Worms Habitat | 3 |
| Food Preservative | 3 |
| Ferments | 2 |
| Industrial | 2 |
| Beverage | 1 |
| Cosmetic | 1 |

61.2% of the total plant species richness, 75.3% of the genera, and 81.6% of the families recorded in the vascular flora of the region by Dávila et al. (2002) and Mendez-Larios et al. (2004, 2005). The 1,605 useful plant species reported here represent nearly 23% of the 7,000 useful vascular plant species probably existing in Mexico according to estimates by Caballero (1984) and Caballero et al. (1998). This information indicates that the Tehuacán–Cuicatlán Valley is Mexico's richest area of ethnobotanical knowledge documented up to now. This fact is particularly relevant since this knowledge is found in a relatively small area (nearly 10,000 km$^2$). Such important information is known in part because of the effort of ethnobotanical research invested in the area, but mostly because of the natural biological diversity characterizing the region (Dávila et al. 2002) as well as its cultural richness (Casas et al. 2001) and history (MacNeish 1967).

The diversified pattern of plant uses, in terms of the number of species and the number of uses per species, coincides with the information generally documented in ethnobotanical studies in Mexico (Berlin et al. 1974; Zizumbo-Villarreal and Colunga-García-Marín 1982; Alcorn 1984; Caballero and Mapes 1985; Bye 1993; Benz et al. 1994; Martínez-Alfaro et al. 1995; Casas et al. 1994, 1996, 2001; Caballero et al. 2000; Lira 2004). This diversified pattern has been considered as a part of what Toledo et al. (2003) called the multiple use of natural resources, characteristic of the form of subsistence of Mesoamerican indigenous rural peoples. The analysis of the spatial distribution of the plant resources and their association with particular ecosystems is yet to be performed, but some attempts to analyze data in this direction (Solís 2006), as well as ecological studies conducted in the area demonstrating the occurrence of high β diversity (Osorio-Beristain et al. 1996; Valiente-Banuet et al. 2000), suggest that ecosystems are reservoirs of particular types of resources and that both the diversity of plant resources and ecosystems are relevant to analyze the pattern of multiple uses of resources and environment of human subsistence in the rural areas of the region.

The residual analysis does not generally corroborate the null hypothesis that families with a higher number of useful species would be those having the higher general species richness. The hypothesis was correct for some families, such as Asteraceae, which contributes a high number of medicinal plant species, but this is not the case for most families. This pattern corroborates that humans have directed their attention to groups with particular qualities of plant resources, rather than to the most abundant or diverse elements of nature (Moerman and Estabrook 2003; Hernández-Delgado et al. 2003, 2005; Caballero et al. 2004; Moerman 2005; Canales et al. 2005, 2006).

The Tehuacán–Cuicatlán Valley has a rich and well–documented history of plant management (MacNeish 1967, 1992; Smith 1967). It is one of the areas widely recognized for the studies on prehistory documenting the origins of agriculture in the New World. Its dry conditions were a determinant in the preservation of signs of early processes of interactions between humans and plants. But the present information suggests that

**Table 4.** The most important plant families in the Tehuacán–Cuicatlán Valley, according to their richness of useful species, in the higher categories of use.

| Family | Total species in the Tehuacán –Cuicatlán Valley (from Dávila et al. 1993) | Useful Species | % Useful Flora | Fodder | Food | Firewood | Medicinal | Ornamental | Wood and Construction Materials |
|---|---|---|---|---|---|---|---|---|---|
| Poaceae | 220 | 220 | 100 | 219 | 12 | 1 | 15 | 7 | 4 |
| Asteraceae | 345 | 195 | 57 | 115 | 23 | 9 | 83 | 34 | 1 |
| Fabaceae | 191 | 80 | 42 | 56 | 20 | 5 | 19 | 4 | 3 |
| Cactaceae | 74 | 67 | 91 | 40 | 44 | 16 | 7 | 16 | 9 |
| Mimosaceae | 72 | 61 | 85 | 49 | 31 | 39 | 13 | 5 | 19 |
| Cyperaceae | 42 | 42 | 100 | 42 | 1 | 0 | 4 | 0 | 0 |
| Euphorbiaceae | 106 | 38 | 36 | 16 | 9 | 1 | 19 | 6 | 2 |
| Solanaceae | 76 | 52 | 68 | 16 | 14 | 0 | 19 | 12 | 0 |
| Fagaceae | 25 | 33 | 132 | 9 | 4 | 33 | 5 | 1 | 32 |
| Malvaceae | 56 | 25 | 45 | 16 | 3 | 1 | 9 | 8 | 0 |
| Caesalpinaceae | 34 | 22 | 65 | 17 | 0 | 10 | 6 | 3 | 4 |
| Bromeliaceae | 47 | 24 | 51 | 20 | 2 | 1 | 4 | 5 | 0 |
| Cucurbitaceae | 21 | 20 | 95 | 8 | 12 | 0 | 4 | 1 | 0 |
| Lamiaceae | 95 | 31 | 33 | 11 | 4 | 0 | 20 | 5 | 0 |
| Verbenaceae | 43 | 16 | 37 | 9 | 8 | 0 | 12 | 3 | 0 |
| Crassulaceae | 49 | 30 | 61 | 2 | 0 | 0 | 6 | 26 | 0 |
| Burseraceae | 19 | 20 | 105 | 5 | 0 | 14 | 6 | 0 | 4 |

**Table 5.** Residual values obtained for the most important useful plant families in the Tehuacán–Cuicatlán Valley.

| Family | Total Species | Useful Species | Calculated | Residual |
|---|---|---|---|---|
| Poaceae | 220 | 220 | 105.6498 | 118.3502 |
| Asteraceae | 345 | 195 | 161.0373 | 33.9627 |
| Fabaceae | 191 | 80 | 92.7999 | −12.7999 |
| Cactaceae | 74 | 67 | 40.9572 | 26.0428 |
| Mimosaceae | 72 | 61 | 40.071 | 20.929 |
| Cyperaceae | 42 | 42 | 26.778 | 26.222 |
| Euphorbiaceae | 106 | 38 | 55.1364 | −17.1364 |
| Solanaceae | 76 | 52 | 41.8434 | 10.1566 |
| Fagaceae | 25 | 33 | 19.2453 | 13.7547 |
| Malvaceae | 56 | 25 | 32.9814 | −7.9814 |
| Caesalpinaceae | 34 | 22 | 23.2332 | −1.2332 |
| Bromeliaceae | 47 | 24 | 28.9935 | −4.9935 |
| Cucurbitaceae | 21 | 20 | 17.4729 | 2.5271 |
| Lamiaceae | 95 | 31 | 50.2623 | −19.2623 |
| Verbenaceae | 43 | 16 | 27.2211 | −11.2211 |
| Crassulaceae | 49 | 30 | 29.8797 | 0.1203 |
| Burseraceae | 19 | 20 | 16.5867 | 3.4133 |
| Acanthaceae | 35 | 22 | 23.6763 | −1.6763 |
| Convolvulaceae | 38 | 20 | 25.0056 | −5.0056 |
| Agavaceae | 25 | 17 | 19.2453 | −2.2453 |

the importance of the Tehuacán Valley is not a question of whether there are appropriate environmental conditions for good preservation of archaeological remains; rather, it appears that the signs found are a representation of intense processes occurring in a fascinating region characterized by high biological diversity and culture whose expression is maintained to the present day.

**Table 6.** Residual values obtained for the plant families from the Tehuacán–Cuicatlán Valley in the most important categories of use.

| Family | Fodder | Ornamental | Wood and Construction Materials | Food | Firewood | Medicinal |
|---|---|---|---|---|---|---|
| Poaceae | 137.5486 | −5.6876 | 0.8737 | −5.2285 | −4.571 | −12.4976 |
| Asteraceae | −13.4139 | 17.0499 | 0.6862 | −0.2785 | 6.8165 | 40.4774 |
| Fabaceae | −14.5561 | −7.6987 | −0.7788 | 4.1751 | −1.3569 | −5.0118 |
| Cactaceae | 13.4008 | 8.291 | 2.5887 | 33.8379 | 6.4724 | −2.9484 |
| Mimosaceae | 23.1522 | −2.6408 | 12.5437 | 20.9347 | 29.4182 | 3.292 |
| Cyperaceae | 36.4232 | – | – | −7.6133 | – | −2.102 |
| Euphorbiaceae | −22.6216 | −2.8002 | −3.6913 | −2.7109 | −7.6604 | 5.2052 |
| Solanaceae | −11.3506 | 4.2228 | – | 3.7411 | – | 8.8112 |
| Fagaceae | 0.8101 | −5.0381 | 24.4862 | −3.7905 | 22.1445 | 0.9414 |
| Malvaceae | −3.8366 | 0.9048 | – | −6.2909 | −9.0154 | 1.2152 |
| Caesalpinaceae | 5.4288 | −3.345 | −3.3113 | – | −0.6116 | 0.8596 |
| Bromeliaceae | 3.5447 | −1.7883 | – | −6.8553 | −9.2593 | −2.703 |
| Cucurbitaceae | 1.3129 | −4.9017 | – | 4.4031 | – | 0.4222 |
| Lamiaceae | −23.4889 | −3.4251 | – | −7.1785 | – | 7.5274 |
| Verbenaceae | −5.9525 | −3.6519 | – | −0.6617 | – | 5.7778 |
| Crassulaceae | −15.2067 | 19.1435 | – | – | – | −0.9434 |
| Burseraceae | −0.9357 | – | −3.6488 | – | 2.9819 | 2.6626 |
| Acanthaceae | 4.0531 | −2.3791 | – | – | – | −1.2606 |
| Convolvulaceae | −1.074 | 2.5186 | −6.2213 | −4.4197 | −8.5032 | 2.3788 |
| Agavaceae | −6.1899 | −1.0381 | −5.5138 | 2.2095 | −4.8555 | −2.0586 |

**Table 7.** Examples of plant species with more than three uses in the Tehuacán–Cuicatlán Valley.

| Species | Family | Uses Number | Uses |
|---|---|---|---|
| *Guazuma ulmifolia* | Sterculiaceae | 11 | Fodder, firewood, food, medicinal, wood and construction materials, ornamental, living fence, fibers, handicrafts, ceremonial, shed |
| *Acacia farnesiana* | Mimosaceae | 10 | Fodder, firewood, medicinal, wood and construction materials, ornamental, living fence, handicrafts, soils control, aromatizing |
| *Alnus acuminata* | Betulaceae | 9 | Fodder, firewood, medicinal, wood and construction materials, ornamental, handicrafts, shed, soils control, colorant |
| *Agave atrovirens* | Agavaceae | 8 | Firewood, medicinal, living fence, fibers, alcoholic beverage, stimulant, source of soap, soils control |
| *Bursera morelensis* | Burseraceae | 7 | Fodder, firewood, wood and construction materials, living fence, resins / latex, soils control, colorant |
| *Stenocereus stellatus* | Cactaceae | 7 | Fodder, firewood, food, wood and construction materials, living fence, alcoholic beverage |
| *Nopalea cochenillifera* | Cactaceae | 6 | Fodder, food, medicinal, ornamental, handicrafts, colorant |
| *Fouquieria formosa* | Fouquieriaceae | 5 | Fodder, firewood, medicinal, wood and construction materials, shed |
| *Fraxinus uhdei* | Oleaceae | 5 | Firewood, medicinal, wood and construction materials, ornamental, shed |
| *Pachycereus hollianus* | Cactaceae | 4 | Firewood, food, wood and construction materials, living fence |
| *Polaskia chende* | Cactaceae | 4 | Fodder, food, living fence, soils control |
| *Polaskia chichipe* | Cactaceae | 4 | Fodder, food, living fence, soils control |

The results of this study illustrate that the old interrelationships between the people and plants of the Tehuacán Valley are ongoing. Most of the plant resources are wild and most of them are obtained by gathering. But recent studies reveal that gathering is not a simple, uniform way of harvesting nature (i.e., see the studies by Arellano and Casas (2003) with *Escontria chiotilla*; González-Soberanis and Casas (2004) with *Sideroxylon palmeri*; Blancas (2007) with *Myrtillocactus schenckii*; and Delgado (2008) with *Agave potatorum*). Gathering involves different techniques and is conducted at different intensities according to a number of social, economic, and cultural factors (González-Insuasti and Caballero 2007; Casas et al. 2008). Currently, people undertake strategies and make agreements in their communities to carry out different routines for gathering and extracting plant resources for a number of purposes (Casas et al. 2008). In relation to the risks caused by the extraction and use of plant resources, these processes constitute a route of research particularly relevant to understanding the early processes of plant management and their relation to the origins of plant domestication (Casas et al. 2007).

Of particular relevance is the information that nearly 500 plant species receive some management type (both silvicultural *in situ* management and/or cultivation). Some studies have documented that these practices may cause modifications in frequencies of phenotypes in the populations of some species and, in some cases, an increase in the frequency of desired phenotypes, as well as morphological differentiation and changes in physiology and life cycles of the managed populations and wild populations (Casas et al. 2007). Accordingly, this group of plants deserves deeper ecological and evolutionary study since the ethnobotanical studies have documented only collaterally the occurrence of such processes.

**Table 8.** Examples of plant species submitted to one or more management forms in the Tehuacán–Cuicatlán Valley.

| Family | Species | Management Forms |
|---|---|---|
| Agavaceae | *Agave peacockii* | Tolerated, cultivated |
| Agavaceae | *Agave potatorum* | Tolerated, cultivated, enhanced |
| Asteraceae | *Montanoa tomentosa* | Tolerated |
| Cactaceae | *Myrtillocactus schenckii* | Tolerated, cultivated |
| Cactaceae | *Myrtillocactus geometrizans* | Tolerated, cultivated |
| Cactaceae | *Pachycereus hollianus* | Tolerated, cultivated |
| Cactaceae | *Polaskia chende* | Tolerated, cultivated |
| Cactaceae | *Polaskia chichipe* | Tolerated, cultivated |
| Cactaceae | *Stenocereus stellatus* | Tolerated, protected, cultivated |
| Cactaceae | *Stenocereus pruinosus* | Tolerated, protected, cultivated |
| Cucurbitaceae | *Sicyos parviflorus* | Tolerated, protected |
| Chenopodiaceae | *Chenopodium ambrosioides* | Tolerated, enhanced |
| Fagaceae | *Quercus glaucoides* | Tolerated, protected |
| Julianaceae | *Juliana adstringens* | Tolerated, protected |
| Malvaceae | *Anoda cristata* | Tolerated, transplanted |
| Mimosaceae | *Leucaena esculenta* | Tolerated, protected, transplanted |
| Sapotaceae | *Sideroxylon palmeri* | Tolerated, cultivated |

Management processes are ongoing, and the plant species that are managed are the expression of resources with a particular cultural value on which people are practicing forms of procurement. Recent studies have documented that some of these management practices have had some effect on the populations of various species, which can be interpreted as forms of incipient domestication. Among these species are the columnar cacti *Escontria chiotilla, Pachycereus hollianus, Polaskia chende, P. chichipe, Stenocereus stellatus*, *S. pruinosus* (Casas et al. 1997b, 1999b, 1999c; Cruz and Casas 2002; Arellano and Casas 2003; Otero-Arnaíz et al. 2003, 2005a, 2005b; Carmona and Casas 2005; Tinoco et al. 2005; Oaxaca-Villa et al. 2006; Rodríguez-Arévalo et al. 2006; Ruíz-Duran 2007), as well as species of other plant groups such as *Sideroxylon palmeri* (Sapotaceae) (González-Soberanis and Casas 2004), *Ceiba aesculifolia* ssp. *parvifolia* (Bombacaceae) (Avendaño et al. 2006), and *Proboscidea louisianica* spp. *fragans* (Pedaliaceae) (Paredes-Flores 2006). In all these studies, the plant uses, their variations, and other aspects related to their biology have been documented and analyzed within the context of domestication by means of the *in situ* management of populations. These examples of *in situ* management represent a particular management intensity of plant resources, in which domestication appears to be occurring, and therefore their study could help in understanding more about processes leading to the origins of agriculture and domestication of plants in a symbolic region for this type of study.

This information, however, is not only important from a theoretical point of view. Such high richness of knowledge and management techniques can be particularly valuable for developing strategies of sustainable management of plant resources. The Tehuacán Valley is a biosphere reserve, and people inhabiting the area live with the particular problems of poverty and scarcity of opportunities to develop their traditional production systems (Valiente-Banuet et al. 2006). The authorities of the reserve, the local people, and the scientists are now in the process of constructing a management plan of the protected area, including the core and buffer zones. The information accumulated through our studies is undoubtedly crucial for recovering traditional practices and designing new ones before the also–new challenges of sustainable forms of using natural resources will take place.

## Acknowledgements

The authors thank the SEMARNAT–CONACYT fund (project C01–0544), DGAPA, UNAM (projects IN220005, IN219608, IN213402, IN210506, IX247904), and the Royal Botanic Gardens, Kew (MGU/Useful Plants Project Mexico), for financial support in

the different steps of the work. We specially thank the people from the Tehuacán–Cuicatlán Valley for allowing us to work in their land.

## Literature Cited

Alcorn, J. 1984. Huastec Mayan Ethnobotany. University of Texas Press, Austin, Texas.

Arellano, E. and A. Casas. 2003. Morphological Variation and Domestication of *Escontria chiotilla* (Cactaceae) under Silvicultural Management in the Tehuacán Valley, Central Mexico. Genetic Resources and Crop Evolution 50:439–453.

Arizmendi, M. C. and A. Espinoza de los Monteros. 1996. Avifauna de los bosques de cactáceas columnares del Valle de Tehuacán, Puebla. Acta Zoológica Mexicana (Nueva Serie) 67:25–46.

Avendaño, A., A. Casas, P. Dávila, and R. Lira. 2006. Use Forms, Management and Commercialization of "Pochote" *Ceiba aesculifolia* (H.B. & K.) Britten & Baker f. subsp. *parvifolia* (Rose) P.E. Gibbs & Semir (Bombacaceae) in the Tehuacán Valley, Central México. Journal of Arid Environments 67:15–35.

Benz, B. F., F. J. Santana, M. R. Pineda, J. Cevallos, L. Robles, and D. De Níz. 1994. Characterization of Mestizo Plant Use in the Sierra de Manantlán, Jalisco–Colima, México. Journal of Ethnobiology 14:23–41.

Berlin, B., D. E. Breedlove, and P. H. Raven. 1974. Principles of Tzeltal Plant Classification: An Introduction to the Botanical Ethnography of a Mayan Speaking Community in Highland Chiapas. Academic Press, New York.

Blancas, J. J. 2007. Manejo tradicional y variación morfológica de *Myrtillocactus schenkii* (J. A. Purpus) Britton & Rose, en el Valle de Tehuacán, Puebla. M. Sc. Thesis. Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México, Mexico.

Blanckaert, I., R. Swennen, M. Paredes-Flores, R. Rosas-López, and R. Lira. 2004. Floristic Composition, Plant Uses and Management Practices in Home Gardens of San Rafael Coxcatlan, Puebla, Mexico. Journal of Arid Environments 57:39–62.

Brailovsky, H., E. Barrera, C. Mayorga, and G. Ortega-León. 1994. Estadíos ninfales de los Coreidos del Valle de Tehuacán, Puebla (Hemiptera–Heteroptera). 1. *Chelinidae staffilesi, C. tabulata* y *Narinia femorata*. Anales del Instituto de Biología, Universidad Nacional Autónoma de México, Serie Zoología 65:241–264.

———, C. Mayorga, G. Ortega-León, and E. Barrera. 1995. Estadíos ninfales de los Coreoidos del Valle de Tehuacán, Puebla, México (Hemiptera–Heteroptera) II. Especies asociadas a huizacheras (*Acacia* spp.) y mezquiteras (*Prosopis* spp.). *Mozena lunata, Pachylis hector, Savius jurgiosus jurgiosus* y *Thasus gigas*. Anales del Instituto de Biología, Universidad Nacional Autónoma de México, Serie Zoología 66:57–80.

Bye, R. A. 1993. The Role of Humans in the Diversification of Plants in Mexico. Pages 707–731 in T. P. Ramamoorthy, R. Bye, A. Lot, and J. Fa, eds., Biological Diversity of Mexico, Origins and Distribution. Oxford University Press, New York.

Caballero, J. 1984. Recursos comestibles potenciales. Pages 114–125 in T. T. Reyna, ed., Seminario sobre la alimentación en México. Instituto de Geografía, Universidad Nacional Autónoma de México, Mexico.

——— and C. Mapes. 1985. Gathering and Subsistence Patterns among the Purhepecha Indians of Mexico. Journal of Ethnobiology 5:31–47.

———, A. Casas, L. Cortés, and C. Mapes. 1998. Patrones en el conocimiento, uso y manejo de plantas en pueblos de México. Estudios Atacameños 16:1–15.

———, A. Casas, L. Cortés, and C. Mapes. 2000. Patrones en el conocimiento, uso y manejo de plantas en pueblos indígenas de México. Revista de Estudios Atacameños 16: 13–31.

———, L. Cortés, M. A. Martínez-Alfaro, and R. Lira. 2004. Uso y Manejo Tradicional de la Diversidad Vegetal. Pages 541–564 in A. García-Mendoza, M. J. Ordoñez, and M. Briones-Salas, eds., Biodiversidad de Oaxaca. Instituto de Biología, UNAM, Fondo Oaxaqueño para la Conservación de la Naturaleza, Mexico–World Wildlife Fund.

Canales, M., T. Hernández, J. Caballero, A. Romo De Vivar, G. Avila, A. Durán, and R. Lira. 2005. Informant Consensus Factor and Antibacterial Activity of the Medicine Plants Used by the People of the San Rafael Coxcatlán, Puebla, México. Journal of Ethnopharmacology 97:429–439.

———, T. Hernández, J. Caballero, A. Romo De Vivar, A. Durán, and R. Lira. 2006. Análisis cuantitativo del conocimiento tradicional de las

plantas medicinales en San Rafael, Coxcatlán, Valle de Tehuacán–Cuicatlán, Puebla, México. Acta Botánica Mexicana 75:21–43.

Carmona, A. and A. Casas. 2005. Management, Phenotypic Patterns and Domestication of *Polaskia chichipe* (Cactaceae) in the Tehuacán Valley, Central Mexico. Journal of Arid Environments 60:115–132.

Casas, A. and J. Caballero. 1996. Traditional Management and Morphological Variation in *Leucaena esculenta* (Fabaceae: Mimosoideae) in the Mixtec Region of Guerrero, Mexico. Economic Botany 50:167–181.

——— and A. Valiente-Banuet. 1995. Etnias, recursos genéticos y desarrollo sustentable en zonas áridas y semiáridas de México. Pages 37–66 in M. Anaya and F. Díaz, eds., IV curso sobre desertificación y desarrollo sustentable en América Latina y el Caribe. Red de Formación Ambiental para América Latina y el Caribe (United Nations Environmental Program)/Red de Cooperación Técnica en Zonas Aridas y Semiáridas de América Latina (Food and Agriculture Organization)/Colegio de Postgraduados en Ciencias Agrícolas, Chapingo, México.

———, J. L. Viveros, and J. Caballero. 1994. Etnobotánica Mixteca: Sociedad, recursos naturales y subsistencia en la Montaña de Guerrero. Consejo Nacional Para la Cultura y las Artes/Instituto Nacional Indigenista, Mexico City.

———, M. C. Vázquez, J. L. Viveros, and J. Caballero. 1996. Plant Management among the Nahua and the Mixtec from the Balsas River Basin: An Ethnobotanical Approach to the Study of Plant Domestication. Human Ecology 24:455–478.

———, J. Caballero, C. Mapes, and S. Zárate. 1997a. Manejo de la vegetación, domesticación de plantas y origen de la agricultura en Mesoamérica. Boletín de la Sociedad Botánica de México 61:31–47.

———, B. Pickersgill, J. Caballero, and A. Valiente-Banuet. 1997b. Ethnobotany and Domestication in Xoconochtli *Stenocereus stellatus* (Cactaceae) in the Tehuacán Valley and La Mixteca Baja, Mexico. Economic Botany 51:279–292.

———, J. Caballero, and A. Valiente-Banuet. 1999a. Use, Management, and Domestication of Columnar Cacti in South–Central Mexico: A Historical Perspective. Journal of Ethnobiology 19:71–95.

———, J. Caballero, A. Valiente-Banuet, J. A. Soriano, and P. Dávila. 1999b. Morphological Variation and the Process of Domestication of *Stenocereus stellatus* (Cactaceae) in Central Mexico. American Journal of Botany 86:522–533.

———, A. Valiente-Banuet, A. Rojas-Martínez, and P. Dávila. 1999c. Reproductive Biology and the Process of Domestication of the Columnar Cactus *Stenocereus stellatus* in Central Mexico. American Journal of Botany 86:534–542.

———, A. Valiente-Banuet, J. L. Viveros, J. Caballero, L. Cortés, P. Dávila, R. Lira, and I. Rodríguez-Arévalo. 2001. Plant Resources of the Tehuacán–Cuicatlán Valley, México. Economic Botany 55:129–166.

———, J. Cruse, E. Morales, A. Otero-Arnaiz, and A. Valiente-Banuet. 2006. Maintenance of Phenotypic and Genotypic Diversity of *Stenocereus stellatus* (Cactaceae) by Indigenous People in Central Mexico. Biodiversity and Conservation 15:879–898.

———, A. Otero-Arnaiz, E. Pérez-Negrón, and A. Valiente-Banuet. 2007. *In situ* Management and Domestication of Plants in Mesoamerica. Annals of Botany 100:1101–1115.

——— and F. Parra. 2007. Agrobiodiversidad, parientes silvestres y cultura: LEISA Revista de Agroecología 23:5–8.

———, S. Rangel-Landa, I. Torres-García, E. Pérez-Negrón, L. Solís, F. Parra, A. Delgado, J. J. Blancas, B. Farfán, and A. I. Moreno. 2008. *In situ* Management and Conservation of Plant Resources in the Tehuacán–Cuicatlán Valley, Mexico: An Ethnobotanical and Ecological Perspective. Pages 1–23 in U. P. De Albuquerque and M. Alves-Ramos, eds., Current Topics in Ethnobotany Research. Signpost, Kerala, India.

Cruz, M. and A. Casas. 2002. Reproductive Biology and Morphological Variation of *Polaskia chende* (Cactaceae) under Domestication in Central Mexico. Journal of Arid Environments 51:561–576.

Dávila, P., J. L. Villaseñor, R. Medina, A. Ramírez, A. Salinas, J. Sánchez-Ken, and P. Tenorio. 1993. Listado Florístico del Valle de Tehuacán–Cuicatlán. Listados florísticos VIII. Instituto de Biología, Universidad Nacional Autónoma de México, Mexico.

———, M. C. Arizmendi, A. Valiente-Banuet, J. L. Villaseñor, A. Casas, and R. Lira. 2002.

Biological Diversity in the Tehuacán–Cuicatlán Valley, Mexico. Biodiversity and Conservation 11:421–442.

Delgadillo, C. and R. H. Zander. 1984. The Mosses of the Tehuacán Valley, Mexico, and Notes on Their Distribution. Bryologist 87:319–322.

Delgado, A. 2008. Aprovechamiento y disponibilidad espacial de *Agave potatorum* en el Valle de Tehuacán–Cuicatlán. M. Sc. Thesis. Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México, Mexico.

Echeverría-Ayala, Y. 2003. Ecología de los recursos vegetales en una comunidad Mixteca del Valle de Tehuacán. Bachelor Thesis. Universidad Michoacana de San Nicolas de Hidalgo, Morelia, Michoacán, México.

Flores, J. S. 1999. Etnobotánica de las leguminosas en la península de Yucatán: Uso y manejo entre los Mayas. Ph.D. Dissertation. Facultad de Ciencias, Universidad Nacional Autónoma de México, Mexico.

González-Insuasti, M. S. and J. Caballero. 2007. Managing Plant Resources: How Intensive Can It Be? Human Ecology 35:303–314.

González-Soberanis, C. and A. Casas. 2004. Traditional Management and Domestication of Tempesquistle, *Sideroxylon palmeri* (Sapotaceae) in the Tehuacán Valley, Central Mexico. Journal of Arid Environments 59:245–258.

Hernández-Delgado, T., M. Canales, J. G. Avila, A. Durán, J. Caballero, A. Romo De Vivar, and R. Lira. 2003. Ethnobotany and Antimicrobial Activity of Some Plants Used in Traditional Medicine of Zapotitlán de las Salinas, Puebla (México). Journal of Ethnopharmacology 88:181–188.

———, M. Canales, J. Caballero, A. Durán, and R. Lira. 2005. Análisis cuantitativo del conocimiento tradicional sobre plantas utilizadas para el tratamiento de enfermedades gastrointestinales en Zapotitlán de las Salinas, Puebla, México. Interciencia 30:529–535.

Hernández-Xolocotzi, E. 1993. Aspects of Plant Domestication in Mexico: A Personal View. Pages 733–753 in T. P. Ramamoorthy, R. Bye, A. Lot, and J. Fa, eds., Biological Diversity of Mexico, Origins and Distribution. Oxford University Press, New York.

Lira, R. 2004. Cucurbitaceae de la Península de Yucatán: Taxonomía, Florística y Etnobotánica. Etnoflora Yucatanense 22. Universidad Autónoma de Yucatán / CONACyT, Mérida, Yucatán, México.

——— and A. Casas. 1998. Uso y manejo en *Ibervillea millspaughii* (Cogn.) C. Jeffrey, *Melothria pendula* L. y otras especies silvestres de la familia Cucurbitaceae: Posibles procesos de domesticación incipiente. Boletín de La Sociedad Botánica de México 62:77–89.

——— and J. Caballero. 2002. Ethnobotany of the Wild Mexican Cucurbitaceae. Economic Botany 56:380–398.

MacNeish, R. S. 1967. A Summary of the Subsistence. Pages 290–331 in D. S. Byers, Ed., The Prehistory of the Tehuacán Valley. Vol. 1: Environment and Subsistence. University of Texas Press, Austin, Texas.

———. 1992. The Origins of Agriculture and Settled Life. University of Oklahoma Press, Norman, Oklahoma.

Martínez-Alfaro, M. M. A., V. Evangelista, M. Mendoza, G. Morales, and A. Wong. 1995. Catálogo de plantas útiles de la Sierra Norte de Puebla, México. Cuadernos 27. Instituto de Biología, Universidad Nacional Autónoma de México, Mexico.

Mendez-Larios, I., J. L. Villaseñor, and E. Ortíz. 2004. Las Magnoliophyta endémicas de la porción xerofítica de la provincia florística del Valle de Tehuacán–Cuicatlán, México. Anales del Instituto de Biología, Universidad Nacional Autónoma de México 75:87–104.

———, J. L. Villaseñor, R. Lira, J. J. Morrone, P. Dávila, and E. Ortíz. 2005. Toward the Identification of a Core Zone in the Tehuacán–Cuicatlán Biosphere Reserve (Mexico), Based on Parsimony Analysis of Endemicity of Flowering Plant Species. Interciencia 30:267–274.

Miranda, F. 1948. Datos sobre la vegetación en la Cuenca Alta del Papaloapan. Anales del Instituto de Biología, Universidad Nacional Autónoma de México 19:333–364.

Moerman, D. 2005. "Symbols and Selectivity" Revisited. Journal of Ethnopharmacology 100:40–42.

——— and G. Estabrook. 2003. Native Americans' Choice of Species for Medicinal Use is Dependent on Plant Family: Confirmation with Meta–Significance Analysis. Journal of Ethnopharmacology 87:51–59.

———, R. W. Pemberton, D. Kiefer, and B. Berlin. 1999. A Comparative Analysis of Five Medicinal Floras. Journal of Ethnobiology 19:49–67.

Oaxaca-Villa, B., A. Casas, and A. Valiente-Banuet. 2006. Reproductive Biology in Wild and Silvicultural Managed Populations of *Escontria chiotilla* (Cactaceae) in the Tehuacán Valley, Central Mexico. Genetic Resources and Crop Evolution 53:277–287.

Osorio-Beristain, O., A. Valiente-Banuet, P. Dávila, and R. Medina. 1996. Tipos de vegetación y diversidad beta en el valle de Zapotitlán de las Salinas, Puebla, México. Boletín de la Sociedad Botánica de México 59:35–58.

Otero-Arnaiz, A., A. Casas, M. C. Bartolo, E. Pérez-Negrón, and A. Valiente-Banuet. 2003. Evolution of *Polaskia chichipe* (Cactaceae) under Domestication in the Tehuacán Valley, Central Mexico: Reproductive Biology. American Journal of Botany 90:593–602.

———, A. Casas, J. L. Hamrick, and J. Cruse. 2005a. Genetic Variation and Evolution of *Polaskia chichipe* (Cactaceae) under Domestication in the Tehuacán Valley, Central Mexico Analyzed by Microsatellite Polymorphism. Molecular Ecology 14:1603–1611.

———, A. Casas, and J. L. Hamrick. 2005b. Direct and Indirect Estimates of Gene Flow among Wild and Managed Populations of *Polaskia chichipe*, An Endemic Columnar Cactus in Central Mexico. Molecular Ecology 14:3313–4322.

Paredes-Flores, M. 2006. Manejo, abundancia y variación morfológica del Torito *Proboscidea louisianica* (Mill.) Thell. ssp. *fragans* (Lindl.) Bretting (Pedaliaceae), en Zapotitlán Salinas, Puebla. M. Sc. Thesis. Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México, Mexico.

———, R. Lira, and P. Davila. 2007. Estudio etnobotánico de Zapotitlán Salinas, Puebla. Acta Botanica Mexicana 79:13–61.

Parra, F., N. Pérez-Nasser, R. Lira, D. Pérez-Salicrup, and A. Casas. 2008. Population Genetics and Process of Domestication of *Stenocereus pruinosus* (Cactaceae) in the Tehuacán Valley, México. Journal of Arid Environments 72(11):1997–2010.

Pennington, C. W. 1969. The Tepehuan of Chihuahua: Their Material Culture. University of Utah Press, Salt Lake City.

Pérez-Negrón, E. and A. Casas. 2007. Use, Extraction Rates and Spatial Availability of Plant Resources in the Tehuacán–Cuicatlán Valley, Mexico: The Case of Quiotepec, Oaxaca. Journal of Arid Environments 70:356–379.

Rangel-Landa, S. and R. Lemus. 2002. Aspectos etnobotánicos y ecológicos de los recursos vegetales entre los Ixcatecos de Santa María Ixcatlán, Oaxaca. Bachelor Thesis. Universidad Michoacana de San Nicolas de Hidalgo, Morelia, Michoacán, México.

Ríos-Casanova, L., A. Valiente-Banuet, and V. Rico-Gray. 2004. Las hormigas del Valle de Tehuacán (Hymenoptera: Formicidae): Una comparación con otras zonas áridas de México. Acta Zoológica Mexicana (Nueva Serie) 20:37–54.

Rodríguez-Arévalo, I., A. Casas, J. Campos, and R. Lira. 2006. Uso, manejo y procesos de domesticación de *Pachycereus hollianus* (F.A.C. Weber) Buxb. (Cactaceae), en el Valle de Tehuacán–Cuicatlán, México. Interciencia 31:677–685.

Rojas-Martínez, A. and A. Valiente-Banuet. 1996. Análisis comparativo de la quiropterofauna del Valle de Tehuacán, Puebla, Oaxaca. Acta Zoológica Mexicana (Nueva Serie) 67:1–23.

Rosas-López, R. 2003. Estudio etnobotánico de San Rafael–Coxcatlán. Bachelor Thesis. Facultad de Estudios Superiores Iztacala, Universidad nacional Autónoma de México, Mexico.

Ruíz-Durán, M. E. 2007. Patrones de diversidad genética y procesos de domesticación de *Polaskia chende* (Cactaceae). Bachelor Thesis. Universidad Michoacana de San Nicolas de Hidalgo, Morelia, Michoacán, México.

Rzedowski, J. 1978. Vegetación de México. Editorial Limusa, México City.

———. 1993. Diversity and Origins of the Phanerogamic Flora of Mexico. Pages 129–144 in T. P. Ramamoorthy, R. Bye, A. Lot, and J. Fa, eds., Biological Diversity of Mexico, Origins and Distribution. Oxford University Press, New York.

Smith, C. E. 1965. Flora, Tehuacán Valley. Fieldiana Botany 31:101–143.

———. 1967. Plant Remains. Pages 220–225 in D. S. Byers, ed., The Prehistory of the Tehuacán Valley. Vol. 1: Environment and Subsistence. University of Texas Press, Austin, Texas.

Solís, L. 2006. Etnoecología cuicateca en San Lorenzo Pápalo, Oaxaca. M. Sc. Thesis. Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México, Mexico.

Tinoco, A., A. Casas, R. Luna, and K. Oyama. 2005. Population Genetics of Wild and

Silvicultural Managed Populations of *Escontria chiotilla* in the Tehuacán Valley, Central Mexico. Genetic Resources and Crop Evolution 52:525–538.

Toledo, V. M., A. I. Batis, R. Becerra, E. Martínez and C. H. Ramos. 1995. La selva útil: etnobotánica cuantitativa de los grupos indígenas del trópico húmedo de México. Interciencia 20:177–187.

———, J. Rzedowski, and J. Villalobos. 1997. Regional Overview: Middle America. Pages 97–124 in S. D. Davis, V. H. Heywood, O. Herrera-MacBryde, J. Villa-Lobos, and A. C. Hamilton, eds., Centres of Plant Diversity. A Guide and Strategy for Their Conservation, Vol. 3: The Americas. IUCN Publications Unit, Cambridge, United Kingdom.

———, P. Alarcón, P. Moguel, M. Olivo, E. Leyequien, A. Cabrera, and A. Rodríguez. 2001. El atlas etnoecológico de México y Centro América: fundamentos, métodos y resultados. Etnoecológica 6:7–41.

———, B. Ortiz–Espejel, L. Cortés, P. Moguel, and M. J. Ordoñez. 2003. The Multiple Use of Tropical Forests by Indigenous Peoples in México: A Case of Adaptive Management. Conservation Ecology 7:3. http://www.consecol.org/vol7/iss3/art9/ (27 January 2009).

Torres, I. 2004. Aspectos etnobotánicos y ecológicos de los recursos vegetales de la comunidad de San Luis Atolotitlán, Mucnicipio de Caltepec, Puebla, México. Bachelor Thesis. Universidad Michoacana de San Nicolas de Hidalgo, Morelia, Michoacán, México.

Valiente-Banuet, L. 1991. Patrones de precipitación en el valle semiárido de Tehuacán, Puebla, México. Bachelor Thesis, Facultad de Ciencias, Universidad Nacional Autónoma de México, Mexico.

———, A. Casas, A. Alcántara, P. Dávila, N. Flores-Hernández, J. L. Villaseñor, J. Ortega-Ramírez, and J. A. Soriano. 2000. La vegetación del Valle de Tehuacán–Cuicatlán. Boletin de la Sociedad Botánica de México 67:25–74.

———, P. Dávila, A. Casas, M. C. Arizmendi, and J. Ortega. 2006. Diversidad biológica y desarrollo sustentable en la reserva de la biosfera Tehuacán Cuicatlán. Pages 248–277 in K. Oyama and A. Castillo, eds., Manejo, Conservación y Restauración de Recursos Naturales en México. Perspectivas desde la Investigación Científica. Siglo XXI Editores, México City.

Villaseñor, J. L., P. Dávila, and F. Chiang. 1990. Fitogeografía del Valle de Tehuacán–Cuicatlán. Boletin de la Sociedad Botánica de México 50:135–149.

Zizumbo-Villarreal, D. and P. Colunga-García-Marín. 1982. Los Huaves: la apropiación de los Recursos Naturales. Universidad Autónoma de Chapingo. Departamento de Sociología Rural.